

*AB* (methylsulfonyl)phenyl]-3-phenyl), Merck L-752,860 and compounds of Formula I--

Please replace the paragraph beginning on page 8, line 26 and ending on page 9, line 5 with the following:

*AB* --Preferred leukotriene B<sub>4</sub> receptor antagonists include Bayer Bay-x-1005 ((R)-α-Cyclopentyl-4-(2-quinolinylmethoxy) benzene acetic acid), Ciba-Geigy CGS-25019C (Benzamide, 4-[[5-[4-(aminoiminomethyl)phenoxy]pentyl]oxy]-3-methoxy-N,N-bis(1-methylethyl)-, (2Z)-2-butenedioate), ebselen (1,2-Benzisoselenazol-3(2H)-one, 2-phenyl), Leo Denmark ETH-615 (Benzoic acid, 4-[[[(3-fluorophenyl)methyl][4-(2-quinolinylmethoxy)phenyl]amino]methyl]), Lilly LY-293111 (Benzoic acid, 2-[3-[3-[(5-ethyl-4'-fluoro-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]), Ono ONO-4057 (Benzenepropanoic acid, 2-(4-carboxybutoxy)-6-[[[(5E)-6-(4-methoxyphenyl)-5-hexenyl]oxy]), Terumo TMK-688 (Carbonic acid, 4-[5-[[2-[4-(diphenylmethoxy)-1-piperidinyl]ethyl]amino]-5-oxo-1,3-pentadienyl]-2-methoxyphenyl ethyl ester), Boehringer Ingelheim BI-RM-270 (2-Benzoxazolamine, N-[(1S)-2-cyclohexyl-1-(2-pyridinyl)ethyl]-5-methyl), Lilly LY 213024 (Benzenepropanoic acid, 5-(3-carboxybenzoyl)-2-(decyloxy)), Lilly LY 264086 (9H-Xanthene-4-propanoic acid, 7-carboxy-3-(decyloxy)-9-oxo), Lilly LY 292728, Ono ONO LB457 (Benzenepropanoic acid, 2-(4-carboxybutoxy)-6-[[[(5E)-6-(4-methoxyphenyl)-5-hexenyl]oxy]), Pfizer 105696, Perdue Frederick PF 10042 (Pyrrolidine, 1-[5-hydroxy-5-[8-(1-hydroxy-2-phenylethyl)-2-dibenzofuranyl]-1-oxopentyl]), Rhone-Poulenc Rorer RP 66153 (2-Thiopheneheptanoic acid, .alpha.,.alpha.-dimethyl-3-(3-phenylpropyl)), SmithKline Beecham SB-201146 (2-Propenoic acid, 3-[6-[[3-(aminophenyl)sulfinyl]methyl]-3-[[8-(4-methoxyphenyl)octyl]oxy]-2-pyridinyl]-, (2E)), SmithKline Beecham SB-201993 (Benzoic acid, 3-[[[[6-[(1E)-2-carboxyethenyl]-5-[[8-(4-methoxyphenyl)octyl]oxy]-2-pyridinyl]methyl]thio]methyl]), Searle SC-53228 (2H-1-Benzopyran-2-propanoic acid, 7-[3-[2-(cyclopropylmethyl)-3-methoxy-4-

[(methylamino)carbonyl]phenoxy]propoxy]-3,4-dihydro-8-propyl-,  
(2S)), Sumitamo SM 15178 (.beta.-Alanine, N-[[6-[(4-acetyl-2-ethyl-5-hydroxyphenoxy)methyl]-2-pyridinyl]carbonyl]-N-ethyl),  
American Home Products WAY 121006 ([1,1'-Biphenyl]-4-acetic acid,  
2-fluoro-4'-(2-quinolinylmethoxy)), Bayer Bay-o-8276, calcitriol  
(9,10-Secocholesta-5,7,10(19)-triene-1,3,25-triol,  
(1.alpha.,3.beta.,5Z,7E)), Warner-Lambert CI-987 (2,4-  
Thiazolidinedione,5-[[3,5-bis(1,1-dimethylethyl)-4-  
hydroxyphenyl]methylene]), Merck and Co. L-651392 (3H-  
Phenothiazin-3-one, 4-bromo-2,7-dimethoxy), Lilly LY 210073,  
Lilly LY 223982 (Benzenepropanoic acid, 5-(3-carboxybenzoyl)-2-  
[[5E)-6-(4-methoxyphenyl)-5-hexenyl]oxy]), Lilly LY-233569 (2-  
Propenamide, N-hydroxy-N-methyl-3-[2-(methylthio)phenyl]), Lilly  
LY-255283 (Ethanone, 1-[5-ethyl-2-hydroxy-4-[[6-methyl-6-(1H-  
tetrazol-5-yl)heptyl]oxy]phenyl]), Merck and Co. MK-591 (1H-  
Indole-2-propanoic acid, 1-[(4-chlorophenyl)methyl]-3-[(1,1-  
dimethylethyl)thio]-a,a-dimethyl-5-(2-quinolinylmethoxy)-, sodium  
salt), Merck and CO. MK-886 (1H-indole-2-propanoic acid, 1-[(4-  
chlorophenyl)methyl]-3-[(1,1-dimethylethyl)thio]-a,a-dimethyl-5-  
(1-methylethyl)), Ono ONO-LB-448, Purdue Frederick PF-5901  
(Benzenemethanol, a-pentyl-3-(2-quinolinylmethoxy)), Rhone-  
Poulenc Rorer RG 14893 (2-Naphthalenecarboxylic acid, 4-[2-  
[methyl(2-phenylethyl)amino]-2-oxoethyl]-8-(phenylmethoxy)),  
Rhone-Poulenc Rorer RP 66364, Rhone-Poulenc Rorer RP 69698  
(Pyridine, 2-[[5-methyl-5-(1H-tetrazol-5-yl)hexyl]oxy]-4,6-  
diphenyl), Searle SC-41930 (2H-1-Benzopyran-2-carboxylic acid, 7-  
[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-  
propyl), Searle SC-50505, Searle SC-51146, SmithKline Beecham  
SK&F-104493 (5H-Pyrrolo[1,2-a]imidazole, 6,7-dihydro-2-(4-  
methoxyphenyl)-3-(4-pyridinyl)), and Teinjin TEI-1338 (Benzoic  
acid, 2-[[4-[2-[2-(2-naphthalenyl)ethenyl]cyclopropyl]-1-  
oxobutyl]amino]-, methyl ester, [1R-[1.alpha.,2.beta.(E)]]).--

Please replace the paragraph on page 9, lines 6-15 with the following:

--More preferred leukotriene B<sub>4</sub> receptor antagonists include Bayer Bay-x-1005 ((R)-a-Cyclopentyl-4-(2-quinolinylmethoxy)benzeneacetic acid), Ciba-Geigy CGS-25019C (Benzamide, 4-[[5-[4-(aminoiminomethyl)phenoxy]pentyl]oxy]-3-methoxy-N,N-bis(1-methylethyl)-, (2Z)-2-butenedioate), ebselen (1,2-Benzisoselenazol-3(2H)-one, 2-phenyl), Leo Denmark ETH-615 (Benzoic acid, 4-[[[(3-fluorophenyl)methyl][4-(2-quinolinylmethoxy)phenyl]amino]methyl]), Lilly LY-293111 (Benzoic acid, 2-[3-[3-[(5-ethyl-4'-fluoro-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]), Ono ONO-4057 (Benzenepropanoic acid, 2-(4-carboxybutoxy)-6-[[5E)-6-(4-methoxyphenyl)-5-hexenyl]oxy]), Terumo TMK-688 (Carbonic acid, 4-[5-[[2-[4-(diphenylmethoxy)-1-piperidinyl]ethyl]amino]-5-oxo-1,3-pentadienyl]-2-methoxyphenyl ethyl ester), Boehringer Ingleheim BI-RM-270 (2-Benzoxazoline, N-[(1S)-2-cyclohexyl-1-(2-pyridinyl)ethyl]-5-methyl), Lilly LY 213024 (Benzenepropanoic acid, 5-(3-carboxybenzoyl)-2-(decyloxy)), Lilly LY 264086 (9H-Xanthene-4-propanoic acid, 7-carboxy-3-(decyloxy)-9-oxo), Lilly LY 292728, Ono ONO LB457 (Benzenepropanoic acid, 2-(4-carboxybutoxy)-6-[[5E)-6-(4-methoxyphenyl)-5-hexenyl]oxy]), Pfizer 105696, Perdue Frederick PF 10042 (Pyrrolidine, 1-[5-hydroxy-5-[8-(1-hydroxy-2-phenylethyl)-2-dibenzofuranyl]-1-oxopentyl]), Rhone-Poulenc Rorer RP 66153 (2-Thiopheneheptanoic acid, .alpha.,.alpha.-dimethyl-3-(3-phenylpropyl)), SmithKline Beecham SB-201146 (2-Propenoic acid, 3-[6-[[3-(aminophenyl)sulfinyl]methyl]-3-[[8-(4-methoxyphenyl)octyl]oxy]-2-pyridinyl]-, (2E)), SmithKline Beecham SB-201993 (Benzoic acid, 3-[[[6-[(1E)-2-carboxyethenyl]-5-[[8-(4-methoxyphenyl)octyl]oxy]-2-pyridinyl]methyl]thio]methyl]), Searle SC-53228 (2H-1-Benzopyran-2-propanoic acid, 7-[3-[2-(cyclopropylmethyl)-3-methoxy-4-[(methylamino)carbonyl]phenoxy]propoxy]-3,4-dihydro-8-propyl-, (2S)), Sumitomo SM 15178 (.beta.-Alanine, N-[[6-[(4-acetyl-2-ethyl-5-hydroxyphenoxy)methyl]-2-pyridinyl]carbonyl]-N-ethyl),

84 and American Home Products WAY 121006 ([1,1'-Biphenyl]-4-acetic acid, 2-fluoro-4'-(2-quinolinylmethoxy)) ---

Please replace the paragraph on page 9, lines 16-19 with the following:

2036000.010302  
--Even more preferred leukotriene B<sub>4</sub> receptor antagonists include Bayer Bay-x-1005 ((R)-α-Cyclopentyl-4-(2-quinolinylmethoxy)benzeneacetic acid), Ciba-Geigy CGS-25019C (Benzamide, 4-[[5-[4-(aminoiminomethyl)phenoxy]pentyl]oxy]-3-methoxy-N,N-bis(1-methylethyl)-, (2Z)-2-butenedioate), ebselen (1,2-Benzisoselenazol-3(2H)-one, 2-phenyl), Leo Denmark ETH-615 (Benzoic acid, 4-[[[(3-fluorophenyl)methyl][4-(2-quinolinylmethoxy)phenyl]amino]methyl]), Lilly LY-293111 (Benzoic acid, 2-[3-[3-[(5-ethyl-4'-fluoro-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]), Ono ONO-4057 (Benzenepropanoic acid, 2-(4-carboxybutoxy)-6-[[[(5E)-6-(4-methoxyphenyl)-5-hexenyl]oxy]), and Terumo TMK-688 (Carbonic acid, 4-[5-[[2-[4-(diphenylmethoxy)-1-piperidinyl]ethyl]amino]-5-oxo-1,3-pentadienyl]-2-methoxyphenyl ethyl ester) ---

IN THE CLAIMS:

Please replace Claim 2 with the following:

84  
2. (once amended) A combination comprising a therapeutically-effective amount of a leukotriene B<sub>4</sub> receptor antagonist and a cyclooxygenase-2 inhibitor selected from Taisho NS-398 (Methanesulfonamide, N-[2-(cyclohexyloxy)-4-nitrophenyl]), meloxicam (2H-1,2-Benzothiazine-3-carboxamide, 4-hydroxy-2-methyl-N-(5-methyl-2-thiazolyl)-, 1,1-dioxide), flosulide (Methanesulfonamide, N-[6-(2,4-difluorophenoxy)-2,3-dihydro-1-oxo-1H-inden-5-yl]), Merck MK-966 (2(5H)-Furanone, 4-[4-(methylsulfonyl)phenyl]-3-phenyl), Merck L-752,860 and compounds of Formula I

2036000.010302